

Research Article

Stomach cancer: epidemiological, diagnostic and therapeutic aspects at the Kara Teaching Hospital, Togo

DOSSOUVI Tamegnon^{1*}, EL-HADJI YAKOUBOU Rafiou²,
ADABRA Komlan³, AMAVI Ayi³, AMOUZOU Efoé-Ga Olivier¹,
KANASSOUA Kokou Kouliwa¹, KASSEGNE Iroukora⁴ and
DOSSEH Ekoué David³

¹Department of General Surgery, CHU-Kara, Togo

²Department of Hepato-Gastroenterology, CHU-Kara, Togo

³Department of General Surgery, CHU, Sylvanus Olympio de Lomé, Togo

⁴Department of General Surgery, CHU-Kara, Tomdè, Togo

Abstract

Objective: To report our experience in the management of gastric cancers at the Kara University Hospital (Togo).

Materials and methods: This was a retrospective and prospective study which was conducted from January 1, 2018, to July 31, 2021, in the general surgery and hepato-gastroenterology department of the Kara University Hospital (Togo). This study involved all patients treated for gastric cancer during the study period at CHU Kara (Togo).

Results: We recorded 32 gastric cancers out of the 218 cases of cancer diagnosed during the study period. We had 20 men and 12 women with a sex ratio (M/F = 1.7). The average age was 58 years with the extremes ranging from 17 to 85 years. The pattern of the consultation was dominated by epigastralgia (100%) and deterioration in general condition (100%). Upper digestive endoscopy with biopsies was performed in all our patients. The antral localization was the most found in 62.5% of cases. The most common macroscopic appearance was ulcers-budding (90.6%) and the dominant histological type was moderately differentiated adenocarcinoma (87.5%). The extension assessment found liver metastasis in 10 cases, multiple regional lymphadenopathies of the hepatic pedicle and celiac trunk in 26 cases, ascites related to peritoneal carcinomatosis in 26 cases. Therapeutically, a 4/5 gastrectomy with D1 dissection was performed in 6 cases; gastrojejunal anastomosis in one case and palliative treatment in 25 cases. Survival at 1 year is 50% (3 patients) among operated patients. All the other patients (78.1%) who received palliative treatment all died within 3 months.

Conclusion: Improving the prognosis of stomach cancer like other cancers requires early diagnosis to perform a gastrectomy, the only guarantee of long survival.

Introduction

Digestive cancers, once rare in Africa, are increasingly occupying an important place in surgical activities [1]. Among these, stomach cancer is one of the most frequent. However, worldwide, the incidence of gastric cancer outside the esophagogastric junction is steadily decreasing by approximately 1.5% per year. This decrease is due to better control of risk factors, foremost among which is the eradication of *Helicobacter Pylori*, which halves the risk of gastric cancer [2,3]. The stomach cancer occupied the second

place of digestive cancers in Kara [4] and is clearly increasing in our surgical practice. The management of digestive cancers in our rural context as in most African countries is confronted with the problems of lack of diagnostic and therapeutic means. The diagnosis is often made late at the stage of complications, therefore, making the management palliative.

The objective of this work is to report our experience of the management of stomach cancer at CHU Kara (Togo).

More Information

***Address for Correspondence:**

DOSSOUVI Tamegnon, General Surgeon, CHU KARA, Togo, Email: dboris@ymail.com

Submitted: February 10, 2022

Approved: March 15, 2022

Published: March 16, 2022

How to cite this article: Tamegnon D, Rafiou EY, Komlan A, Ayi A, Olivier AE, et al. Stomach cancer: epidemiological, diagnostic and therapeutic aspects at the Kara Teaching Hospital, Togo. Arch Surg Clin Res. 2022; 6: 001-003.

DOI: 10.29328/journal.ascr.1001062

Copyright: © 2022 Tamegnon D, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Keywords: Stomach cancer; Diagnostic delay; Gastrectomy; Chemotherapy



Materials and methods

This was a retrospective and prospective study which was conducted from January 1, 2018, to June 31, 2021, in the general surgery and hepato-gastroenterology department of the Kara University Hospital (Togo). This study concerned all patients treated for gastric cancer during the study period. The pathological anatomy department is not available at the Kara University Hospital (Togo). The parameters studied were epidemiological (frequency, sex, age, profession, history), clinical, paraclinical, therapeutic, postoperative, and follow-up.

Results

During the study period, we registered 218 cases of digestive cancers, including 32 gastric cancers. We had 20 men and 12 women with a sex ratio (M/F = 1.7). The average age was 58 years with the extremes ranging from 17 to 85 years.

The patient history was characterized by a gastric ulcer in 22 cases and epigastralgia in 10 cases.

The pattern of the consultation was dominated by epigastralgia (100%) and deterioration in general condition (100%) as illustrated in Table 1.

One patient was classified as OMS 1, one OMS 2, and thirty patients OMS 4.

The physical examination revealed a gastric mass in only one case.

Upper digestive endoscopy with biopsies was performed in all our patients. Antral localization was the most found in 62.5% of cases. The most common macroscopic appearance was ulcero-budding (90.6%) and the dominant histological type was moderately differentiated adenocarcinoma (87.5%) as shown in Table 2.

The extension assessment found liver metastasis in 10 cases, multiple regional lymphadenopathies of the hepatic pedicle and celiac trunk in 26 cases, ascites related to peritoneal carcinomatosis in 26 cases. The staging of the cases is done in Table 3.

Therapeutically, a 4/5 gastrectomy with D1 dissection was performed in 6 cases; gastrojejunal bypass in one case and palliative treatment in 25 cases. Adjuvant chemotherapy could not be performed for several reasons: the absence of an oncology service in the Kara region and the poverty of the patients.

The postoperative course was marked by parietal suppuration in one case which regressed under local care. On the digestive level, the operated patients had resumed transit after an average of 3 days and had quickly resumed food, which improved their general condition.

Table 1: Pattern of consultation.

	n	%
Epigastralgia	32	100
Deterioration of general condition	32	100
Dysphagia	7	21.9
Gastrointestinal bleeding	5	15.6

Table 2: Topography, macroscopic and microscopic appearance of gastric cancers.

	n	%
Topography		
Antre	20	62,5
Fundus	5	15,6
Fundus	3	9,4
Diffus	3	9,4
Cardia	1	3,1
Total	32	100
Macroscopic appearance		
Ulcero-budding	29	90,6
Budding	2	6,3
Infiltrating	1	3,1
Histological type		
Adénocarcinoma		
Moderately differentiated	28	87,4
Poorly differentiated	2	6,3
Well differentiated	2	6,3

Table 3: Staging of stomach cancer.

Stage	n	%
Stage 1	0	0
Stage 2	6	18,8
Stage 3	0	0
Stade 4	26	81,2
Total	32	100

Survival at 1 year is 50% (3 patients) among operated patients. All the other patients (78.1%) who received palliative treatment all died within 3 months.

Discussion

Gastric cancer is unevenly distributed around the world. And for good reason, there is a certain genetic and environmental susceptibility to developing this cancer. It is in East Asia that it is highest, especially in Japan and China, followed by Central Eastern and Southern Europe, particularly Portugal and South America [4]. A study of gastric cancers in Africa showed a marked increase in the incidence of this cancer in Mali compared to other African countries such as Uganda and Senegal. Similarly, this incidence is higher in sub-Saharan Africa than in North Africa [5]. These geographic variations may be related to differences in risk factors, such as the prevalence rate of *Helicobacter pylori* infection [6] and accessibility to diagnostic means. Stomach cancer occupies actually the second place in the digestive cancers in CHU Kara [4]. The absence of a cancer registry in Togo and the unavailability of global data (only hospital) mean that the real incidence of this pathology is underestimated.

The average age in our series is 58 years old. Stomach cancer is rare before the age of 40, qualifying it as that of the



elderly. However, this cancer is increasingly encountered in young subjects [7,8] as in our study. The male predominance of this cancer has been mentioned in several studies [7,8].

Several factors favoring the occurrence of this cancer have been incriminated, the first of which is *Helicobacter Pylori*, which is acquired since early childhood [9,10]. In our series, the gastric ulcer was the most common contributing factor in 68.8% of cases. This would be due to an infection with *Helicobacter Pylori* which could not be sought for lack of diagnostic means. Apart from this bacterium having carcinogenic properties, other factors such as tobacco, alcohol, excessive consumption of salt, smoking, curing, canning still occupy an important part in the diet. The artisanal activities of smoking and salting fish are widely established in the coastal regions of West Africa and their impact on the development of gastric cancers is still little known [10].

After upper endoscopy, the antral site (62.5%) was by far the most frequent. This remark has been the rule in the literature contrasting with the rarity of cardia localization [9,10].

The circumstances of the discovery of these cancers in our series were represented essentially by epigastralgia (100%) and deterioration of the general state (100%). The deterioration of the general state is a sign of poor prognosis testifying to the advanced stage of cancers diagnosed in our context like the majority of African countries [8-10]. This is explained by a delay in consultation due to ignorance, attachment to traditional therapy, and the poverty of the African [10]. The other digestive cancers as in the case of liver cancers which are sometimes transferred lesions of gastric origin (31.3%), representing the most frequent digestive cancers in our context, do not escape these diagnostic problems late [11].

Therapeutically, the recommended neo and adjuvant chemotherapy could not be performed in our series due to the high cost not being accessible to patients and the unavailability of chemotherapy in our region. This would have improved the prognosis [12].

The tumor resection rate is very low in our series due to late-stage diagnosis like other digestive cancers.

The one-year survival time in our series is very short due to the absence of complementary treatment. This survival was better in the series by Diop, et al. in Dakar where the majority of patients received adjuvant chemotherapy [10].

Postoperative follow-up was difficult in our work context where patients are struggling to meet appointments and honor additional examinations.

Conclusion

Stomach cancer is one of the most common cancers in the Kara region. Gastric ulcer is the most found contributing factor. The diagnosis was made mainly at the late stage, which made gastrectomy impossible, which is the only curative treatment to improve survival.

References

1. ALLIAM. Les cancers en Afrique francophone. La Ligue Nationale contre le Cancer (France). 2017.
2. Chapelle N, Manfredi S, Lepage C, Faivre J, Bouvier AM, et al. Trends in gastric cancer incidence: a period and birth cohort analysis in a well-defined French population. *Gastric Cancer*. 2016; 19: 508–514. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/26118904/>
3. Rokkas T. A systematic review and meta-analysis of the role of *Helicobacter pylori* eradication in preventing gastric cancer. *Ann Gastroenterol*. 2017; 30: 414-423. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/28655977/>
4. Dossouvi T, Yakoubou RE, Bouglouga O, Kanassoua KK, Kassegne I, et al. Profil épidémiologique des cancers digestifs au chu-kara (Togo). *J Afr Chir Digest*. 2021; 21: 3464-3468.
5. Asombang AW, Kelly P. Gastric cancer in Africa: what do we know about incidence and risk factors? *Trans R Soc Trop Med Hyg*. 2012; 106: 69-74. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/22136952/>
6. Wong BC, Lam SK, Ching CK, Hu WH, Kwok E, et al. Differential *Helicobacter pylori* infection rates in two contrasting gastric cancer risk regions of South China. *J Gastroenterol Hepatol*. 1999; 14: 120–125. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/10029291/>
7. Fadlouallah M, Krami H, Errabih I, Benzoubeir N, et al. Le cancer gastrique : aspects épidémiologiques au Maroc. *J Afr Cancer*. 2014; 1-7.
8. Diarra MT, Konaté A, Diarra AN, Sow HEC, Doumbia KES, et al. Epidemiological characteristics and prognosis of gastric Cancer in urban areas of Mali. *Mali Med*. 2014; 29: 50-54. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/30049116/>
9. Bouglouga O, Lawson-Ananissouh LM, Bagny A, Kaaga L, Amegbor K. Stomach cancer: Epidemiological, clinical and histological aspects at the Lome Campus teaching hospital (Togo). *Med Sante Trop*. 2015; 25: 65-85. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/25786857/>
10. Diop B, Dia AA, Ba PA, Sow O, Thiam O, et al. Prise en charge chirurgicale des tumeurs gastriques à Dakar: à propos de 36 observations. *Health Sci. Dis*. 2017; 18: 34-38.
11. Bouglouga O, Bagny A, Lawson-Ananissouh LM, Djibril MA, Redah D, et al. La prise en charge du carcinome hépatocellulaire progresse-t-elle en Afrique noire? *Rev Med Adag*. 2012; 2: 176-179.
12. Russo AE, Strong VE. Gastric Cancer Etiology and Management in Asia and the West. *Ann Rev Med*. 2018; 70: 353–367.